

Comment on "Experimental observation of the topological structure of exceptional points" [Phys.Rev.Letters 86, 787 (2001)]

I. Rotter

Max-Planck-Institut für Physik komplexer Systeme, D-01187 Dresden, Germany
(February 1, 2008)

Dembowski et al. [1] report on a microwave cavity experiment where exceptional points (EP) are encircled. The authors claim that the EP is a fourth order branch point for the wave functions because the two wave functions are not just interchanged as their corresponding energies during a complete loop in the parameter (λ) plane, but one of them undergoes a phase change. Therefore, the EP can, according to the authors, be clearly distinguished from other topological singularities such as diabolic points (DP). The last ones are encircled some years ago in a similar experiment [2]. In the following, it will be shown that the two experiments [1] and [2] prove the same topological singularity in contrast to the statement of [1].

The encircled EP [1,3] is nothing else than a second order branch point in the complex plane (BPCP) with the following properties [4]:

- (i) the complex energies of the two crossing states are equal, $E_1 - \frac{i}{2}\Gamma_1 = E_2 - \frac{i}{2}\Gamma_2$,
- (ii) their wave functions are linearly dependent,

$$\psi_1 = \pm i \psi_2, \quad (1)$$

and bi-orthogonal with $\langle \psi_i | \psi_i \rangle \rightarrow \infty$, $|\langle \psi_i | \psi_{i \neq j} \rangle| \rightarrow \infty$ and $\langle \psi_i^* | \psi_j \rangle = \delta_{i,j}$. Although the number of BPCP is of measure zero, their influence on the level dynamics is quite large. It can be traced to, e.g., the well-known avoided crossing of discrete or resonance states [4]. The BPCP and DP are therefore related to one another.

For illustration, let us consider the complex two-by-two Hamiltonian matrix

$$\mathcal{H} = \begin{pmatrix} e_1(\lambda) - \frac{i}{2}\gamma_1 & \omega \\ \omega & e_2(\lambda) - \frac{i}{2}\gamma_2 \end{pmatrix} \quad (2)$$

where e_k and γ_k ($k = 1, 2$) are the unperturbed energies and widths, respectively, of the two states. The e_k are assumed to depend on the parameter λ in such a manner that the two states may cross in energy at λ^{cr} when $\omega = 0$. The two states interact only via ω which is assumed in the following to be independent of the parameter λ (as the γ_k). The eigenvalues of $\mathcal{H}$ are $E_{\pm} - \frac{i}{2}\Gamma_{\pm} = (e_1 + e_2) - \frac{i}{2}(\gamma_1 + \gamma_2) \pm \frac{1}{2}\sqrt{F}$ with

$$F = ((e_1 - e_2) - \frac{i}{2}(\gamma_1 - \gamma_2))^2 + 4\omega^2. \quad (3)$$

When $F(\lambda, \omega) = 0$ at $\lambda = \lambda^{\text{cr}}$ (and $\omega = \omega^{\text{cr}}$), the S matrix has a double pole which is a second order BPCP.

According to (3), $F = F_R + i F_I$ is generally a complex number. For real ω , $e_1 = e_2$ at $\lambda = \lambda^{\text{cr}}$ and we have to differentiate between three cases

$$F_R(\lambda, \omega) > 0 \rightarrow \sqrt{F_R} = \text{real} \quad (4)$$

$$F_R(\lambda, \omega) = 0 \rightarrow \sqrt{F_R} = 0 \quad (5)$$

$$F_R(\lambda, \omega) < 0 \rightarrow \sqrt{F_R} = \text{imag}. \quad (6)$$

The first case gives the avoided level crossing in energy with an exchange of the two wave functions at λ^{cr} . The second case corresponds to the double pole of the S matrix. In the third case, the two levels cross freely in energy and the two states are *not* exchanged at the critical value λ^{cr} [4]. In [5], the two cases $F_R > 0$ and $F_R < 0$ are studied experimentally in a microwave cavity and called *overcritical* and *subcritical* coupling, respectively.

The DP is surrounded in the experiment [2] in the regime of overcritical coupling along the whole way of encircling and λ^{cr} is passed twice in opposite directions, (i) $\psi_1 \rightarrow -i\psi_2$, $\psi_2 \rightarrow +i\psi_1$ and (ii) $-i\psi_2 \rightarrow -\psi_1$, $i\psi_1 \rightarrow -\psi_2$. The phase change occurring after surrounding the DP corresponds to the geometric phase discussed by Berry [6].

The way of encircling the BPCP itself passes from a region with overcritical coupling at λ^{cr} to another one with subcritical coupling at λ^{cr} . Thus, a first full surrounding gives $\psi_1 \rightarrow -i\psi_2$, $\psi_2 \rightarrow +i\psi_1$ and a second one (in the same direction) $-i\psi_2 \rightarrow +\psi_1$, $i\psi_1 \rightarrow +\psi_2$. That means, surrounding the BPCP twice restores the wave functions ψ_i including their phases. The BPCP is a second order branch point in correspondence with the eigenvalue equation [4]. Encircling the BPCP in the opposite direction gives $\psi_1 \rightarrow +i\psi_2$, $\psi_2 \rightarrow -i\psi_1$ in agreement with the experimental data of [1].

Summarizing, the phase changes observed by encircling a DP and a BPCP (called EP in [1,3]) are caused by the *same* topological singularity.

-
- [1] C. Dembowski, H.D. Gräf, H.L. Harney, A. Heine, W.D. Heiss, H. Rehfeld and A. Richter, Phys. Rev. Lett. **86**, 787 (2001)
 - [2] H.M. Lauber, P. Weidenhammer and D. Dubbers, Phys. Rev. Lett. **72**, 1004 (1994)
 - [3] W.D. Heiss, M. Müller, and I. Rotter, Phys. Rev. E **58**, 2894 (1998)
 - [4] I. Rotter, arXiv:quant-ph/0105068 and quant-ph/0107018
 - [5] M. Philipp, P. von Brentano, G. Pascovici, and A. Richter, Phys. Rev. E **62**, 1922 (2000)
 - [6] M.V. Berry, Proc. R. Soc. London A **392**, 45 (1984)